



Summary

The handbook of multidisciplinary and integrated heart failure care



September 2018

The Heart Failure Policy Network is an independent, multidisciplinary platform made possible with financial support from Novartis Pharma. The content produced by the Network is not biased to any specific treatment or therapy, and is endorsed and owned by the Network's members, who have full editorial control. All members provide their time for free.

Executive summary

Heart failure

#1

cause of unplanned admissions

1 in 5

can expect to live with HF at some point in their lives

+50%

rise in hospital admissions due in the next 25 years

Best-practice care achieves

↓30%

reduction in hospitalisations

Heart failure (HF) is a healthcare sustainability challenge.

- **HF is a common condition.** It occurs as a result of the heart becoming too weak or stiff,¹ affecting its ability to pump enough blood around the body.²
- **At least 15 million people live with HF in Europe.**³ One in five of us can expect to live with HF at some point in our lives.⁴
- **The burden of HF is high.**^{5,6} Quality of life and survival remain poor – worse than for most common types of cancer.⁷
- **HF is the leading cause of unplanned hospital readmissions.**⁸ It is also the most common cause of admissions in people over 65.⁹
- **The burden of HF will rise.** This is partly due to an ageing population and improved survival from other cardiovascular and chronic diseases.^{10,11}
- **Hospital admissions due to HF are projected to rise by 50%** over the next 25 years.⁹



Although the prognosis is poor, the right package of care makes a huge difference to people living with HF.

- **The right care and support can allow people with HF to recover many years of life,** and quality of life.^{12,13} Hospitalisation can also be reduced by up to 30%.¹⁴⁻¹⁷
- **Care and support must be flexible to the individual needs and preferences** of the person living with HF. This can improve clinical outcomes and patient activation to live and self-manage HF.^{2,6,18}
- **The best model of care is an HF management programme:** a package of person-centred care, which includes self-management support, rehabilitative and preventive care, routine reviews and escalation in the event of crisis.^{6,19}
- **Best practice is for multidisciplinary care to be led by specialists,** including cardiologists (ideally with a sub-specialty in HF) and HF specialist nurses, working out of HF clinics.^{6,12} Other vital roles include GPs, cardiac rehabilitation specialists, physiotherapists and pharmacists.
- **Home visits and structured telephone support (led by HF specialist nurses) are innovative models** in reducing HF-specific admissions and mortality.¹⁷

European health systems are currently unprepared for HF.

- **Healthcare systems often struggle with a chronic disease model,** and HF is no exception. Care is often fragmented^{6,19} and guideline-based care too rare.^{2,9,20}
- **The greatest challenges and missed opportunities lie at five points across the HF journey:** presentation and diagnosis; hospital discharge and follow-up; clinical management; patient empowerment and self-care; and advance care planning.
- **We need to train new roles and reinforce existing ones** if we are to keep patients out of hospital. In particular, we lack HF specialist nurses,¹⁹ and need to better train and involve GPs and pharmacists. Every healthcare professional should recognise basic symptoms.
- **We need to overcome inertia, low awareness and low scrutiny at all levels,** including among policymakers, government agencies, professionals, patients and the public.

Call to action

We call on governments to recognise heart failure (HF) as an urgent sustainability challenge for 21st-century healthcare systems. Governments across Europe must:

- **Have a formal strategy on HF** and the changing impact it will have on the healthcare system and society, including future scenario-modelling. This should be developed in close consultation with patient and clinical advocates.
- **Invest in sustainable, specialist HF care models** outside of acute care, for example HF specialist nurses and HF outpatient centres. They should also promote professional education and, where appropriate, additional specialist accreditation for GPs, internists, primary care nurses and expert patients.
- **Ensure national guidelines and local care pathways embed the vision of quality in routine delivery,** working with professional societies, patient advocacy groups and healthcare providers to do so.
- **Prepare robust and public national audits of performance** to ensure accountability to citizens on patient survival, quality of life and experience of care, and to guide investment and incentives. The safe reduction of hospital readmissions should be a major strategic goal, indicating a sustainable approach to HF care.

We call on governments to demonstrate measurable improvement for the following minimum standards and core indicators of quality for all HF patients:

- **Specialist-led diagnosis.** It is vital to achieve a definitive diagnosis with an echocardiogram, ensuring the underlying causes of HF are fully understood, addressed and communicated to patients.
- **Natriuretic peptide testing.** This should be a routinely available tool in both primary and secondary care settings.
- **Specialist-led care in hospital.** Patients should receive input from a cardiologist and HF specialist nurse.
- **Hospital discharge with a care plan.** Discharge plans should include clear points of contact and timely follow-up by specialists.
- **Cardiac rehabilitation, patient therapeutic education and psychological support.** HF care must be built on the maximum engagement of patients.
- **A shift in management of HF from the acute to primary care setting.** This shift should occur wherever it is safe and effective to do so.

The following organisations support and endorse this handbook.



The handbook of multidisciplinary and integrated heart failure care and supplementary material are available at www.hfpolicynetwork.eu

References

- NHS Choices. 2016. Heart Failure: Overview. Available from: <https://www.nhs.uk/conditions/heart-failure/> [Accessed 20/08/18]
- Ponikowski P, Anker SD, AlHabib KF, et al. 2014. Heart failure: preventing disease and death worldwide. *ESC Heart Fail* 1(1): 4-25
- Dickstein K, Cohen-Solal A, Filippatos G, et al. 2008. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2008. *European Journal of Heart Failure* 10(10): 933-89
- Lloyd-Jones DM, Larson MG, Leip EP, et al. 2002. Lifetime risk for developing congestive heart failure: the Framingham Heart Study. *Circulation* 106(24): 3069-72
- Jeon YH, Kraus SG, Jowsey T, et al. 2010. The experience of living with chronic heart failure: a narrative review of qualitative studies. *BMC Health Serv Res* 10: 77
- Ponikowski P, Voors A, Anker S, et al. 2016. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC). *European Heart Journal* 37(27): 2129-200
- Mamas MA, Sperrin M, Watson MC, et al. 2017. Do patients have worse outcomes in heart failure than in cancer? A primary care-based cohort study with 10-year follow-up in Scotland. *Eur J Heart Fail* 19(9): 1095-104
- Gheorghide M, Vaduganathan M, Fonarow GC, et al. 2013. Rehospitalization for Heart Failure: Problems and Perspectives. *Journal of the American College of Cardiology* 61(4): 391-403
- Cowie M, Anker S, Cleland J, et al. 2014. Improving care for patients with acute heart failure: before, during and after hospitalization. *ESC Heart Failure* 1(2): 110-45
- National Institute for Health and Care Excellence. 2010. *Chronic heart failure in adults: management*. London: NICE
- Frankenstein L, Fröhlich H, Cleland J. 2015. Multidisciplinary approach for patients hospitalized with heart failure. *Revista Española de Cardiología (English edition)* 68(10): 885-91
- McDonagh T, Blue L, Clark A, et al. 2011. European Society of Cardiology Heart Failure Association Standards for delivering heart failure care. *European Journal of Heart Failure* 13(3): 235-41
- National Institute for Health and Care Excellence. 2018. *Chronic heart failure: Management of chronic heart failure in adults in primary and secondary care (update)*. Draft for consultation. London: NICE
- McAlister F, Stewart S, Ferrua S, et al. 2004. Multidisciplinary strategies for the management of heart failure patients at high risk for admission: a systematic review of randomized trials. *J Am Coll Cardiol* 44(4): 810-9
- Comin-Colet J, Verdu-Rotellar J, Vela E, et al. 2014. Efficacy of an integrated hospital-primary care program for heart failure: a population-based analysis of 56,742 patients. *Revista Española de Cardiología (English edition)* 67(4): 283-93
- Comin-Colet J, Enjuanes C, Verdu-Rotellar JM, et al. 2016. Impact on clinical events and healthcare costs of adding telemedicine to multidisciplinary disease management programmes for heart failure: Results of a randomized controlled trial. *J Telemed Telecare* 22(5): 282-95
- Feltner C, Jones C, Cene C, et al. 2014. Transitional care interventions to prevent readmissions for persons with heart failure: a systematic review and meta-analysis. *Ann Intern Med* 160(11): 774-84
- Ekman I, Wolf A, Olsson LE, et al. 2012. Effects of person-centred care in patients with chronic heart failure: the PCC-HF study. *Eur Heart J* 33(9): 1112-9
- Glogowska M, Simmonds R, McLachlan S, et al. 2015. Managing patients with heart failure: a qualitative study of multidisciplinary teams with specialist heart failure nurses. *Ann Fam Med* 13(5): 466-71
- Cleland JGF, Swedberg K, Follath F, et al. 2003. The EuroHeart Failure survey programme—a survey on the quality of care among patients with heart failure in Europe Part 1: patient characteristics and diagnosis. *European Heart Journal* 24(5): 442-63
- NHS England. Integrated care and support [online]. Available from: <https://www.england.nhs.uk/ourwork/part-rel/transformation-fund/> [Accessed 26/07/18]
- Jaarsma T, Stromberg A. 2014. Heart failure clinics are still useful (more than ever?). *Can J Cardiol* 30(3): 272-5
- Scottish Intercollegiate Guidelines Network. 2016. *Management of chronic heart failure*. Edinburgh: SIGN
- Bradley EH, Sipsma H, Horwitz LI, et al. 2015. Hospital strategy uptake and reductions in unplanned readmission rates for patients with heart failure: a prospective study. *J Gen Intern Med* 30(5): 605-11

The Heart Failure Policy Network is an independent, multidisciplinary platform made possible with financial support from Novartis Pharma. The content produced by the Network is not biased to any specific treatment or therapy, and is endorsed and owned by the Network's members, who have full editorial control. All members provide their time for free.

What is multidisciplinary and integrated care in heart failure?

Multidisciplinary and integrated care is about delivering the best care possible

Integrated care has been defined as realising the potential of multidisciplinary teams to promote person-centred and coordinated care, tailored to the needs and preferences of the patient, their family and carers.²¹ Effective care of HF should ensure management of comorbidities, changing needs and support throughout different care settings, involving HF specialists.⁶ Team members are expected to work in close coordination with one another – including the patient – with mutual respect, clear communication and clear division of responsibilities.¹⁹ This is especially important when a person with HF transitions between care settings.

Multidisciplinary and integrated care changes across the HF journey

Every person living with HF has a unique journey. However, there are three very typical phases of HF: diagnosis; care and follow-up; and living with HF.^{2,22} The last phase represents the long-term experience of people who learn to live with the condition and self-care. Depending on progression of the disease, people with HF may have episodes that require return to acute care or enhanced medical supervision.

The handbook of multidisciplinary and integrated heart failure care was developed by the Heart Failure Policy Network Secretariat in conjunction with the 2018 Project Advisory Group, which comprises patient advocates, cardiologists and representatives from professional associations. The Heart Failure Policy Network would like to thank and acknowledge their contribution as well as that of all members of the Network and case study leads who provided their time and insights.



The handbook of multidisciplinary and integrated heart failure care and supplementary material are available at www.hfpolicynetwork.eu

Figure 1. Multidisciplinary and integrated heart failure care journey: key components of quality

Seamless transitions of care are required across moments of crisis and changing need, in all settings and stages of the patient journey.^{2,6}

